# Hand and arm transplants

Clinical Policy ID: CCP.1321

Recent review date: 9/2026

Next review date: 1/2028

Policy contains: Composite tissue allotransplantation; forearm transplant; hand transplant.

*First Choice Next has developed clinical policies to assist with making coverage determinations. First Choice Next's clinical policies are based on guidelines from established industry sources, such as the Centers for Medicare & Medicaid Services (CMS), state regulatory agencies, the American Medical Association (AMA), medical specialty professional societies, and peer-reviewed professional literature. These clinical policies along with other sources, such as plan benefits and state and federal laws and regulatory requirements, including any state- or plan-specific definition of "medically necessary," and the specific facts of the particular situation are considered, on a case by case basis, by First Choice Next when making coverage determinations. In the event of conflict between this clinical policy and plan benefits and/or state or federal laws and/or regulatory requirements, the plan benefits and/or state and federal laws and/or regulatory requirements shall control. First Choice Next's clinical policies are for informational purposes only and not intended as medical advice or to direct treatment. Physicians and other health care providers are solely responsible for the treatment decisions for their patients. First Choice Next's clinical policies are reflective of evidence-based medicine at the time of review. As medical science evolves, First Choice Next will update its clinical policies as necessary. First Choice Next's clinical policies are not guarantees of payment.*

## Coverage policy

Hand and arm transplants are investigational/not clinically proven and, therefore, not medically necessary.

Limitations

All other uses of hand and arm transplants are not medically necessary.

Alternative covered services

- Limb replantation.
- Upper limb prosthesis.
- Physical and occupational therapy.

## Background

Upper-limb amputations make up a minority – between one fifth and one sixth – of total amputations. Seventy percent of upper-limb amputations are below the elbow, with 10% of these at the hand or wrist (Fahrenkopf, 2018).

Composite tissue allotransplantation is the transfer of vascularized or non-vascularized heterogeneous tissues with different antigenicities from one person to another. Unlike a solid organ transplant, vascularized composite tissue allotransplantation involves multiple tissues (e.g., skin, muscle, tendon, bone, cartilage, fat, nerves, and blood vessels). Composite tissue allotransplantation is a complex procedure requiring multidisciplinary technical skills and understanding of its immunologic aspects. It is a treatment for complex injuries that leave patients with

structural, functional, and aesthetic deficits that cannot be addressed by other means (American Society of Transplantation, 2015).

Hand and arm transplantations are forms of vascularized composite tissue allotransplantation. Hand transplantation is an extremely complex procedure that can last from eight to ten hours; following the procedure, transplant recipients require life-long immunosuppressive therapy and intensive physical therapy to regain hand and arm function (The Johns Hopkins University, 2024).

The first hand transplant was attempted in 1964 in South America without success. In the United States, advances in surgical technique and immunosuppression allowed for the first successful hand transplant in 1999. Vascularized composite tissue allotransplantation is regulated, as are all solid organ transplants, within the Organ Procurement and Transplantation Network (80 FR 26464). Since federal oversight began in July 2014, 70 candidates have been registered for vascularized composite tissue transplants other than uterus, and 37 have undergone transplantation through 2024. Upper limb was the most frequently performed type, accounting for 16 of the 37 procedures. Nationally, 21 programs have performed at least one such transplant, of which nine have performed an upper limb transplant. In 2024, three procedures were performed, one of them upper limb (Nichols, 2026).

## Findings

The evidence on upper extremity transplantation consistently highlights an uncertain risk-benefit profile, a conclusion informing a cautious approach across all evidence types. Clinical guidelines, particularly from health technology assessment bodies, question the procedure's value due to uncertain long-term benefits and high costs (Health Quality Ontario, 2016; National Institute for Health and Care Excellence, 2011). This stance is supported by systematic reviews that, while noting potential functional gains (Mordukhovich, 2026; Wells, 2022), quantitatively detail significant risks, including high rates of acute rejection and severe complications from lifelong immunosuppression (Huelsboemer, 2024b; Knoedler, 2026; Milek, 2023). Other evidence from clinical reports and comparative studies further underscores this caution, establishing stringent patient selection criteria (Mendenhall, 2020) and situating the procedure as one of several options, including advanced prosthetics (Efanov, 2022).

Guidelines:

Official guidelines from major professional and health technology assessment bodies frame hand transplantation as a specialized therapeutic option rather than a standard procedure, citing an uncertain balance of benefits and risks. This is exemplified by Health Quality Ontario, which recommends against publicly funding the procedure due to both uncertain clinical benefit and poor value for money (Health Quality Ontario, 2016). Similarly, the National Institute for Health and Care Excellence states that current evidence on the efficacy and safety of the procedure is “inadequate in quantity,” while the American Society for Surgery of the Hand and the Transplantation Society of Australia and New Zealand both emphasize that it is a quality-of-life procedure that may introduce life-shortening risks from lifelong immunosuppression (American Society for Surgery of the Hand, 2013; Dwyer, 2019; National Institute for Health and Care Excellence, 2011). Consequently, guidelines impose stringent patient selection criteria; for example, the Transplantation Society of Australia and New Zealand endorses the procedure only for bilateral amputees and encourages a trial with prosthetics first, a more restrictive approach than in other jurisdictions (Dwyer, 2019). Ultimately, all guidance documents mandate that the procedure be performed exclusively in specialized centers with multidisciplinary teams experienced in both hand surgery and transplant medicine, and under special arrangements for clinical governance, consent, and ongoing research to address the current evidence gaps (American Society for Surgery of the Hand, 2013; Health Quality Ontario, 2016; National Institute for Health and Care Excellence, 2011). In 2025, the National Academies of Sciences, Engineering, and Medicine issued a report on protocols and standard operating procedures for face

and hand transplants, recommending a whole-health approach delivered at high-quality transplant centers, standardization of care management and data collection across the field, and a structured framework for ongoing research. The Organ Procurement and Transplantation Network committee reviewed and supported these recommendations (Nichols, 2026).

Systematic Reviews

Complications and Immunosuppression

Systematic reviews report that acute rejection affects a majority of recipients, with rates ranging from 65.9% in a hand transplant-only cohort to over 90% in a cohort of mixed transplant types (Huelsboemer, 2024b; Milek, 2023). A broader analysis across 211 vascularized composite allograft cases found an average of 1.45 acute rejections per patient, with higher rates in burn patients than trauma patients (3.73 vs 1.78 average rejections) (Van Dieren, 2024). These events contribute to the risk of graft failure, with one review reporting graft loss in 13.2% of hand transplant patients and another citing allograft removal in 22% of its cohort (Huelsboemer, 2024b; Milek, 2023).

These complications are managed with intensive immunosuppressive regimens that carry their own significant risks. The common protocol, used in over 90% of hand transplant patients, involves induction therapy followed by triple maintenance therapy with tacrolimus, mycophenolate mofetil, and steroids (Huelsboemer, 2024b; Van Dieren, 2024). This long-term therapy is associated with high rates of opportunistic infections, impaired glucose metabolism (62%), and renal insufficiency (26%) (Milek, 2023). This has led some authors to conclude that the side effects of immunosuppression "remain difficult to reconcile" with the benefits of a procedure that is not life-saving (Huelsboemer, 2024b).

A systematic review of 46 studies, encompassing (n = 136) recipients of vascularized composite tissue, of whom (n = 69) received upper extremity transplants, examined treatment of rejection. Among upper extremity recipients, rejection treatment preserved the graft in 54 cases and was unsuccessful in nine, four of which followed discontinuation of immunosuppressive therapy because of infection or unrelated surgery. Corticosteroids were the mainstay of treatment, given alone in 22 cases and combined with topical tacrolimus in 10; treatment duration ranged from two days to three months. The authors reported that chronic rejection still lacks an accepted definition, a staging system, and a validated treatment algorithm, and is likely under-recognized rather than rare (Knoedler, 2026).

To better manage these risks, research has focused on identifying biomarkers of rejection. A systematic review that analyzed data from 22 articles, encompassing (n = 26) patients, investigated this issue. The review found that donor specific antibodies were significantly associated with higher rejection grades (mean grades of 2.7 versus 2.2, P=0.005). The authors noted, however, that the utility of their findings was limited by inconsistent reporting between centers and the absence of standardized diagnostic guidelines for antibody-mediated rejection (Huelsboemer, 2024a).

Functional Outcomes and Patient Selection

Evidence from systematic reviews suggests that functional improvements are achievable, though the certainty of these findings is limited. An earlier review found that disability scores of the arm, shoulder, and hand declined significantly, indicating improvement, following transplantation (Wells, 2022). However, the literature is characterized by a lack of standardized assessment tools and significant heterogeneity in outcome measures, making direct comparison across studies difficult. Achieving positive outcomes is also linked to patient selection; a review of practices at transplantation centers identified the patient's capacity to manage the allograft post-transplantation—including access to follow-up and psychological stability—as a key criterion (Laspro, 2023).

A systematic review of 118 studies, encompassing (n = 97) patients who received 148 transplanted limbs between 1998 and 2025 across 14 countries, examined the timeline of recovery over the first 10 postoperative

years. Level of transplantation influenced outcome: power grip strength was greater following distal than proximal transplantation (median 7.6 kilograms versus 4.5 kilograms, P<.01). Total active range of motion of the fingers did not change significantly after the first postoperative year in either group. Bilateral recipients reported better general health status on the 36-Item Short Form Health Survey than unilateral recipients (median 74.5 versus 53.5, P<.05), although disability scores of the arm, shoulder, and hand did not differ significantly between the two groups. Thirteen limbs were reamputated across 12 patients, at a median of 2.3 years after surgery. The authors noted that most included records were low level of evidence and that outcome measures remain heterogeneous across centers (Mordukhovich, 2026).

A systematic review and meta-analysis of 20 survey studies examined public willingness to donate and to receive vascularized composite tissue. For hand transplantation, pooled willingness to donate was 54.8% (95% confidence interval 45.7% to 63.8%), drawn from (n = 11,172) participants, and pooled willingness to receive a hand transplant was 72.4% (95% confidence interval 65.1% to 79.7%). Four studies directly compared attitudes toward solid organ and vascularized composite tissue donation, and all four found respondents significantly more willing to donate solid organs. Three surveys found that between 18% and 54% of respondents would be less likely to register as organ donors if vascularized composite tissue were automatically included in donor registration (Olivas, 2026).

Other Evidence

Patient selection criteria are further detailed in reports based on clinical experience. These sources frequently cite an age range of 18 to 69 years, an absence of significant medical or psychosocial issues, and no recent history of cancer as important factors for candidacy (MacKey, 2014; Mendenhall, 2020). A qualitative study of 50 candidates and recipients confirmed that patient preferences for candidacy were similar to these established criteria (Vanterpool, 2023). An important consideration in this selection process is the comparison to alternatives; a comparative study of 26 hand transplants and 45 myoelectric prostheses found no significant difference in overall quality-adjusted life-years (P=.36), but did find significantly higher quality-adjusted life-years for prostheses in the subgroup of unilateral amputees (P=.0015) (Efanov, 2022).

In 2025, the findings were revised thematically and included previously uncited guidelines (American Society for Surgery of the Hand, 2013; Dwyer, 2019) and recent systematic reviews (Huelsboemer, 2024a, 2024b; Van Dieren, 2024). No policy changes were warranted.

In 2026, the findings were updated with recent systematic reviews of functional recovery (Mordukhovich, 2026), rejection treatment (Knoedler, 2026), and public perceptions of donation (Olivas, 2026), along with current national registry data (Nichols, 2026). No policy changes were warranted.

## References

On August 15, 2026, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were “Vascularized Composite Allotransplantation” (MeSH), “Transplantation, Homologous" (MeSH), “Upper Extremity” (MeSH), “Composite Tissue Allografts” (MeSH), “arm transplant,” and “hand transplant.” We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.

80 FR 26464. Organ Procurement and Transplantation: Implementation of the HIV Organ Policy Equity Act.

American Society for Surgery of the Hand Council. ASSH position statement on hand transplantation 2013. *J Hand Surg Am.* 2013; 38(11):2234-5. Doi:10.1016/j.jhsa. 2013.08.119.

American Society of Transplantation. Vascularized composite allotransplantation (VCA) research. https://www.myast.org/public-policy/key-position-statements/vascularized-composite-allotransplantation-vca-research. Revised August 11, 2015.

Dwyer, K. M., Burt, J. D., & Bennett, T. International vascularised composite allotransplantation activity: implications for Australia. Med J Aust. 2019; 210 (2). Doi: 10.5694/mja2.12068.Efanov JI, Izadpanah A, Bou-Merhi J, Lin SJ, Danino MA. Applying health utility outcome measures and quality-adjusted life-years to compare hand allotransplantation and myoelectric prostheses for upper extremity amputations. *Plast Reconstr Surg*. 2022;149(3):465e-474e. Doi: 10.1097/PRS.0000000000008846.

Fahrenkopf MP, Adams NS, Kelpin JP, Do VH. Hand amputations. *Eplasty*. 2018;18:ic21. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6173827/.

Health Quality Ontario. Composite tissue transplant of hand or arm: OHTAC recommendation. https://www.hqontario.ca/evidence-to-improve-care/health-technology-assessment/reviews-and-recommendations/composite-tissue-transplant-of-hand-or-arm. Published June 2016.

Heineman J, Bueno EM, Kiwanuka H, et al. All hands on deck: Hand replantation versus transplantation. *SAGE Open Med.* 2020;8:2050312120926351. Doi: 10.1177/2050312120926351.

Huelsboemer L, Moscarelli J, Dony A, Boroumand S, Kochen A, et al. The role of C4d and donor-specific antibodies in face and hand transplantation—a systematic review in Vascularized Composite Allotransplantation. Frontiers in Transplantation. 2024 a;3:1442006. Doi:10.3389/frtra.2024.1442006.

Huelsboemer L, Moscarelli J, Dony A, Boroumand S, Kochen A, et al. Immunosuppressive strategies in face and hand transplantation: a comprehensive systematic review of current therapy regimens and outcomes. Frontiers in Transplantation. 2024 b;3:1366243. Doi:10.3389/frtra.2024.1366243.

Honeyman C, Stark HL, Fries CA, et al. Vascularised composite allotransplantation in solid organ transplant recipients: A systematic review. *J Plast Reconstr Aesthet Surg*. 2021;74(2):316-326. Doi: 10.1016/j.bjps.2020.08.052.

Knoedler L, Niederegger T, Schaschinger T, et al. A systematic review of treatment strategies to combat acute and chronic rejection episodes in vascularized composite allotransplantation. Frontiers in Immunology. 2026;17:1733221. Doi: 10.3389/fimmu.2026.1733221.

Laspro M, Onuh OC, Carrion K, et al. Vascularized composite allotransplantation of the hand: A systematic review of eligibility criteria. *Ann Plast Surg*. 2023;91(6):771-778. Doi: 10.1097/sap.0000000000003659.

MacKay BJ, Nacke E, Posner M. Hand transplantation – A review. *Bull Hosp Jt Dis.* 2014;72(1):76-88. http://hjdbulletin.org/files/archive/pdfs/8.pdf.

Mendenhall SD, Brown S, Ben-Amotz O, Neumeister MW, Levin LS. Building a hand and upper extremity transplantation program: Lessons learned from the first 20 years of vascularized composite allotransplantation. *Hand (NY)*. 2020;15(2):224-233. Doi: 10.1177/1558944718790579.

Milek D, Reed LT, Echternacht SR, et al. A systematic review of the reported complications related to facial and upper extremity vascularized composite allotransplantation. *J Surg Res*. 2023;281:164-175. Doi: 10.1016/j.jss.2022.08.023.

Mordukhovich I, Roncagli C, Hoffner N, et al. Timeline of functional recovery following hand and upper-extremity transplantation. J Hand Surg Am. Published online May 9, 2026:1.e1-1.e14. Doi: 10.1016/j.jhsa.2026.05.003.

National Institute for Health and Care Excellence. Hand allotransplantation. Interventional procedures guidance [IPG383]. https://www.nice.org.uk/guidance/ipg383. Published March 2011.

Nichols DS, Testa G, Johannesson L, et al. OPTN/SRTR 2024 annual data report: Vascularized composite allograft. Am J Transplant. 2026;26(2 Suppl 1):S585-S612.

Olivas CG, Wyatt HP, Gursky AK, et al. Global public perceptions of vascularized composite allotransplantation: A systematic review. Ann Plast Surg. 2026;97(3):355-366. Doi: 10.1097/SAP.0000000000004831.

The Johns Hopkins University. Comprehensive Transplant Center. Comprehensive hand transplant program. https://www.hopkinsmedicine.org/transplant/programs/reconstructive-transplant/hand-arm-transplant.html. Published 2024.

Van Dieren L, Tawa P, Coppens M, et al. Acute Rejection Rates in Vascularized Composite Allografts: A Systematic Review of Case Reports. *J Surg Res*. 2024;298:137-148. Doi:10.1016/j.jss.2024.02.019.

Vanterpool KB, Gacki-Smith J, Downey MC, et al. Patient preferences of patient selection criteria for upper extremity vascularized composite allotransplantation: A qualitative study. *SAGE open medicine*. 2023;11:20503121231181236. Doi: 10.1177/20503121231181236.

Wells MW, Rampazzo A, Papay F, Gharb BB. Two decades of hand transplantation: A systematic review of outcomes. *Ann Plast Surg*. 2022;88(3):225-344. Doi: 10.1097/SAP.0000000000003056.

## Policy updates

7/2017: initial review date and clinical policy effective date: 8/2017

9/2018: Policy references updated. Policy ID changed.

9/2019: Policy references updated.

9/2020: Policy references updated.

9/2021: Policy references updated.

9/2022: Policy references updated.

9/2023: Policy references updated.

9/2024: Policy references updated.

9/2025: Policy references updated.

9/2026: Policy references updated.

## Related codes

Below are the most commonly submitted codes for the service(s)/item(s) subject to this policy CCP.1321 This is not an exhaustive list of codes. Providers are expected to consult the appropriate coding manuals and bill accordingly.

| Code | Code description |
|---|---|
| 20802 | Replantation, arm (includes surgical neck of humerus through elbow joint), complete amputation |
| 20805 | Replantation, forearm (includes radius and ulna to radial carpal joint), complete amputation |
| 20808 | Replantation, hand (includes hand through metacarpophalangeal joints), complete amputation |
| 20999 | Unlisted procedure, musculoskeletal system, general |

| Code | Code description |
|---|---|
| S2152 | Solid organ(s), complete or segmental, single organ or combination of organs; deceased or living donor(s), procurement, transplantation, and related complications; including: drugs; supplies; hospitalization with outpatient follow-up; medical/surgical, diagnostic, emergency, and rehabilitative services, and the number of days of pre- and post-transplant care in the global definition |