# Prostatic Urethral Lift/UroLift for Benign Prostatic Hyperplasia

Clinical Policy ID: CCP.1529

Recent review date: 8/2026

Next review date: 12/2027

Policy contains: benign prostatic hyperplasia; prostatic urethral lift; UroLift.

*First Choice Next has developed clinical policies to assist with making coverage determinations. First Choice Next's clinical policies are based on guidelines from established industry sources, such as the Centers for Medicare & Medicaid Services (CMS), state regulatory agencies, the American Medical Association (AMA), medical specialty professional societies, and peer-reviewed professional literature. These clinical policies along with other sources, such as plan benefits and state and federal laws and regulatory requirements, including any state- or plan-specific definition of "medically necessary," and the specific facts of the particular situation are considered, on a case by case basis, by First Choice Next when making coverage determinations. In the event of conflict between this clinical policy and plan benefits and/or state or federal laws and/or regulatory requirements, the plan benefits and/or state and federal laws and/or regulatory requirements shall control. First Choice Next's clinical policies are for informational purposes only and not intended as medical advice or to direct treatment. Physicians and other health care providers are solely responsible for the treatment decisions for their patients. First Choice Next's clinical policies are reflective of evidence-based medicine at the time of review. As medical science evolves, First Choice Next will update its clinical policies as necessary. First Choice Next's clinical policies are not guarantees of payment.*

## Coverage policy

Prostatic urethral lift (UroLift, Teleflex, Inc., Pleasanton, California) is clinically proven and, therefore, may be medically necessary for treatment of lower urinary tract symptoms due to benign prostatic hyperplasia when all of the following criteria are met (American Urological Association [Goueli, 2026]):

- Member has a prostate volume between 30 and 80 cc.
- There is verified absence of an obstructive median lobe of the prostate.

Limitations

For members with a prostate volume greater than 80 cc and up to 100 cc or with an obstructive median lobe, the decision to proceed with a prostatic urethral lift procedure should be made on a case-by-case basis, in appropriate members and by clinicians with sufficient procedural experience/expertise (Goueli, 2026).

Contraindications to UroLift include the following (U.S. Food and Drug Administration, 2025):

- Prostate volume of greater than 100 cc
- A urinary tract infection
- Urethral conditions that may prevent insertion of delivery system into bladder
- Urinary incontinence
- Current gross hematuria
- A known allergy to nickel

Alternative covered services

- Medications, including alpha blockers, 5-alpha reductase inhibitors, or a combination
- Transurethral resection of the prostate
- Guideline-directed minimally invasive surgery, including:
    - Convective radiofrequency water vapor thermal therapy
    - Prostatic arterial embolization
    - Temporary implantable nitinol device
    - Transurethral microwave thermotherapy

## Background

Benign prostatic hyperplasia, also known as benign prostatic hypertrophy, is a nonmalignant growth of prostate tissue. The condition is marked by symptoms of the lower urinary tract, urinary retention, or infections due to incomplete bladder emptying. Some cases will not require treatment but can be addressed by watchful waiting to ensure worsening of symptoms is limited. Other cases can be treated conservatively with alpha blockers, 5-alpha reductase inhibitors, phosphodiesterase-5 inhibitors (tadalafil), antimuscarinics, or a combination. However, these medications are not always effective and are associated with elevated risk of ejaculatory and erectile dysfunction (Ng, 2024).

For cases requiring surgery, transurethral approaches and enucleation procedures have largely replaced open prostatectomy as preferred surgical options. Minimally invasive surgical options such as paclitaxel-coated prostatic balloon dilation, transurethral microwave thermotherapy, water vapor or steam infusion therapy, and prostatic urethral internal lateral suturing (prostatic urethral lift) have emerged, offering shorter operating room time, faster recovery, and fewer side effects (Ng, 2024).

The prostatic urethral lift is an endoscopic procedure that retracts obstructing prostatic lobes using small metal implants to secure the retracted position of the enlarged prostate tissue away from the urethra. A disposable cartridge delivers an implant consisting of a capsular nitinol tab and a urethral stainless steel tab held together by a non-absorbable suture, which draws the prostatic urethra to the capsule. The procedure creates an open channel from the bladder neck to the verumontanum. It requires three to four tabs per implantation and either local or general anesthesia, and it can be performed in inpatient or outpatient settings (Rahman, 2024).

In 2013, the U.S. Food and Drug Administration approved the UroLift System UL400 for the treatment of benign prostatic hyperplasia in patients age 50 years and older with no obstructive median or lateral lobe hyperplasia and prostate volumes between 30 and 80 cc (U.S. Food and Drug Administration, 2013). In 2017, approval was expanded to include the UL500 model for lateral and median lobe prostate hyperplasia (U.S. Food and Drug Administration, 2017).

In 2019, approval for recent models expanded based on substantial equivalence to predicate devices and unpublished data presented to the U.S. Food and Drug Administration. Approval includes individuals with prostate volumes up to 100 cc and patients aged 45 years and older. Lowering the age criterion was based on

an early American Urological Association guideline defining the index patient as older than 45 years with lower urinary tract symptoms and multiple studies demonstrating minimal differences between the populations age 45 and age 50 in terms of histopathology, volume, and symptomatology (U.S. Food and Drug Administration, 2025).

## Findings

Guidelines

The American Board of Urology reports that the use of prostatic urethral lift procedures has increased significantly since its introduction in 2015 and currently account for one-third of all procedures for benign prostatic hyperplasia (Zhang, 2023).

According to the American Urological Association (Goueli, 2026), there are a range of procedural options for benign prostatic hyperplasia that differ in their mechanisms of action, anesthesia requirements, hospital length of stay, bleeding risk, and side effect profiles. Indications for early use of procedural therapies include: renal insufficiency, refractory urinary retention, recurrent bladder stones, gross hematuria; recurrent urinary tract infections, recurrent bladder stones, or gross hematuria secondary to benign prostatic hyperplasia; lower urinary tract symptoms/benign prostatic hyperplasia refractory to other therapies; or unwillingness to use other therapies (Clinical Principle). Clinicians should practice shared decision-making weighing the efficacy and durability of each procedure against individual side effect and retreatment profiles (Clinical Principle).

The American Urological Association guideline recommends prostatic urethral lift for patients with lower urinary tract symptoms from benign prostatic hyperplasia who meet the following criteria (Moderate Recommendation; Evidence Level: Grade B) (Goueli, 2026):

- Prostate volume is 30 to 80 cc and verified absence of an obstructive median lobe.
- For prostate volume up to 100 cc or those with an obstructing median lobe, this procedure should only be considered in appropriate patients and by clinicians with sufficient procedural experience/expertise.

These recommendations were based on the inclusion criteria and results of the L.I.F.T. study (ClinicalTrials.gov identifier NCT01294150). The inclusion criteria were participants ages 50 years and older with an International Prostate Symptom Score > 12, a peak flow rate (Qmax) ≤ 12 mL/s, and a prostate volume 30 to 80 cc. Participants were randomized to the Lift procedure or sham control and followed for five years. Prostatic urethral lift offered rapid improvement in symptoms, quality of life, and flow rate durable to five years with a higher likelihood of preserving sexual function compared to many other surgical interventions (Roehrborn, 2017).

Evidence review

Recent systematic reviews/meta-analyses produced the following findings on the effectiveness of prostatic urethral lift/UroLift. While the prostatic urethral lift improves symptoms from a risk-benefit perspective, it is generally not as effective as transurethral resection of the prostate (Cornu, 2023). Similarly, prostatic urethral lift showed little to no difference in urological symptom improvement compared to transurethral resection of the prostate, although it was the most efficacious among five minimally invasive procedures (Franco, 2021, 2022).

Current evidence supports prostatic urethral lift/UroLift for patients with small prostate volumes ranging from 30 to 80 cc without obstructive median lobes based on the L.I.F.T. randomized, sham-controlled trial. The evidence for those with larger prostate volumes (81 to 100 cc) and with obstructive median lobes is very limited in the published literature. Results of the MedLift prospective, nonrandomized study (n = 45) were insufficient to support prostatic urethral lift as a safe and effective treatment for patients with benign prostatic hyperplasia and obstructive median lobes (Rukstalis, 2019).

Long-term effectiveness and safety

The long-term effectiveness and safety of prostatic urethral lift have been highlighted in several studies. Jing (2020) observed that the effects of prostatic urethral lift weaken over time, with patients tracked up to 24 months, and that while it was not as effective as transurethral resection of the prostate, prostatic urethral lift remained safe and effective in selected patients. Tanneru (2020) supported these findings, reporting that prostatic urethral lift was well-tolerated and provided favorable outcomes in symptoms and sexual health over a 24-month period. Sajan (2022) noted that prostatic urethral lift had similar symptom improvement and adverse event rates compared to other minimally invasive procedures at three, six, and 12 months, but transurethral resection of the prostate consistently yielded superior outcomes during these periods.

A meta-analysis by Xiang (2020) consolidated data from 19 articles, covering 11 independent patient series and a total of 304 to 605 patients. The study found significant improvements in the International Prostate Symptom Score by 9.73 to 12.16 points, the Benign Prostatic Hyperplasia Impact Index by 3.74 to 4.50 points, and the maximum flow rate by 3.44 to 4.26 milliliters per second over 24 months. Quality of life scores also improved by 2.20 to 2.55 points, with stable or slightly improved sexual function. Complications were minimal and typically mild, with no significant changes in postvoid residual volume, supporting prostatic urethral lift as an effective and safe procedure that preserves sexual function.

Re-intervention rates and cost effectiveness

Re-intervention rates and cost effectiveness are critical factors in evaluating the overall utility of prostatic urethral lift. Miller (2020) analyzed data from 11 studies involving 2,016 patients and found a pooled annual surgical re-intervention rate of 6.0%, with variations depending on follow-up duration. This highlights a higher re-intervention rate than commonly cited in the literature, emphasizing the need for long-term follow-up data. Chughtai (2022) noted that prostatic urethral lift had lower improvements in prostate scores than other procedures and the highest five-year cost, approximately $9,580 compared to $6,328 for transurethral resection of the prostate. Despite these costs, Light (2021) found that prostatic urethral lift had the highest rate of erectile function preservation at one, six, 12, and 24 months compared with other minimally invasive procedures.

Comparative effectiveness of prostatic urethral lift and other treatments

Several studies have compared the effectiveness of prostatic urethral lift with other treatments over varying periods. Baboudjian (2023) reported that after five years, the effectiveness of surgical or minimally invasive retreatment was 13% for prostatic urethral lift versus 4% for water vapor thermal therapy. Lucas-Cava (2023) found that prostatic urethral lift had a significantly higher rate of re-interventions but a significantly lower rate of major adverse events compared to transurethral resection of the prostate. Minimally invasive procedures such as prostatic urethral lift did not result in significant changes in ejaculatory or erectile function and was associated with a lower risk of retrograde ejaculation compared to transurethral resection of the prostate, other electrosurgical procedures, and laser treatment (Busetto, 2025; Cahill, 2026; Gemma, 2024; Manfredi, 2022). Page (2021) noted that after prostatic urethral lift, the in-hospital complication rate was 3.4%, with 93% of patients being catheter-free within 30 days, and retreatment rates at one and two years were 5.2% and 11.9%, respectively.

In 2024, we revised the coverage section based on updated clinical guidelines from American Urological Association. We also revised the findings section to group studies thematically and added new systematic reviews (Miller, 2020; van Kollenburg, 2023; Xiang, 2020).

In 2025, we updated the references and revised the medical necessity criteria to align with current American Urological Association guideline recommendations for prostatic urethral lift procedures. These changes include deleting the age criterion and the requirement of medication failure.

In 2026, we updated the references, including the American Urological Association guideline. No major policy changes are warranted.

## References

On June 8, 2026, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were “benign prostatic hyperplasia,” “benign prostatic hypertrophy,” “prostatic urethral lift,” and “UroLift.” We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.


Baboudjian M, Cornu J-N, Gondran-Tellier B, et al. Pharmacologic and surgical retreatment after office-based treatments for benign prostatic hyperplasia: A systematic review. *Eur Urol Focus*. 2023;9(5):727-733. Doi: 10.1016/j.euf.2023.03.004.

Busetto GM, Lombardo R, De Nunzio C, et al. Ejaculation sparing of classic and minimally invasive surgical treatments of LUTS/BPH. *Prostate Cancer Prostatic Dis*. 2025;28(1):37-44. Doi: 10.1038/s41391-024-00834-y.

Cahill EM, Lindenbaum MM, Estright A, et al. The impact of minimally invasive surgical therapy for benign prostatic hyperplasia on sexual function: A systematic review and meta-analysis. *Prostate Cancer Prostatic Dis*. 2026;Doi: 10.1038/s41391-026-01091-x.

Chughtai B, Rojanasarot S, Neeser K, et al. A comprehensive analysis of clinical, quality of life, and cost-effectiveness outcomes of key treatment options for benign prostatic hyperplasia. *PLoS One*. 2022;17(4):e0266824. Doi: 10.1371/journal.pone.0266824.

ClinicalTrials.gov. The safety and effectiveness of UroLift: LIFT Pivotal Study (LIFT). ClinicalTrials.gov identifier NCT01294150. https://www.clinicaltrials.gov/study/NCT01294150?term=NCT01294150&rank=1. Last update posted September 5, 2018.

Cornu J-N, Zantek P, Burtt G, et al. Minimally invasive treatments for benign prostatic obstruction: A systematic review and network meta-analysis. *Eur Urol*. 2023;83(6):534-547. Doi: 10.1016/j.eururo.2023.02.028.

Franco JV, Jung JH, Imamura M, et al. Minimally invasive treatments for lower urinary tract symptoms in men with benign prostatic hyperplasia: A network meta-analysis. *Cochrane Database Syst Rev*. 2021;7(7):CD013656. Doi: 10.1002/14651858.CD013656.pub2.

Franco JVA, Jung JH, Imamura M, et al. Minimally invasive treatments for benign prostatic hyperplasia: A Cochrane network meta-analysis. *BJU Int*. 2022;130(2):142-156. Doi: 10.1111/bju.15653.

Gemma L, Pecoraro A, Sebastianelli A, et al. Impact of minimally invasive surgical procedures for male lower urinary tract symptoms due to benign prostatic hyperplasia on ejaculatory function: A systematic review. *Prostate Cancer Prostatic Dis*. 2024;27(3):404-421. Doi: 10.1038/s41391-024-00795-2.

Goueli R, Badlani GH, Welliver C, et al. Management of lower urinary tract symptoms attributed to benign prostatic hyperplasia: AUA guideline (2026) part iii: Procedural/surgical management. *J Urol*. 2026;216(2):161-170. Doi: 10.1097/ju.0000000000005099.

Jing J, Wu Y, Du M, et al. Urethral lift as a safe and effective procedure for prostatic hyperplasia population: A systematic review and meta-analysis. *Front Surg*. 2020;7:598728. Doi: 10.3389/fsurg.2020.598728.

Light A, Jabarkhyl D, Gilling P, et al. Erectile function following surgery for benign prostatic obstruction: A systematic review and network meta-analysis of randomized controlled trials. *Eur Urol*. 2021;80(2):174-187. Doi: 10.1016/j.eururo.2021.04.012.

Lucas-Cava V, Sanchez-Margallo FM, Gorbea II, Sun F. Comparative efficacy and safety of prostatic urethral lift vs prostatic artery embolization for benign prostatic hyperplasia: A systematic review and network meta-analysis. *BJU Int*. 2023;131(2):139-152. Doi: 10.1111/bju.15748.

Manfredi C, Garcia-Gomez B, Arcaniolo D, et al. Impact of surgery for benign prostatic hyperplasia on sexual function: A systematic review and meta-analysis of erectile function and ejaculatory function. *Eur Urol Focus*. 2022;8(6):1711-1732. Doi: 10.1016/j.euf.2022.06.007.

Miller LE, Chughtai B, Dornbier RA, McVary KT. Surgical reintervention rate after prostatic urethral lift: Systematic review and meta-analysis involving over 2,000 patients. *J Urol*. 2020;204(5):1019-1026. Doi: 10.1097/JU.0000000000001132.

Ng M, Leslie SW, Baradhi KM. Benign prostatic hyperplasia. In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2026 Jan-. https://www.ncbi.nlm.nih.gov/books/NBK558920/. Updated October 20, 2024.

Page T, Veeratterapillay R, Keltie K, Burn J, Sims A. Prostatic urethral lift (UroLift): A real-world analysis of outcomes using hospital episodes statistics. *BMC Urol*. 2021;21(1):55. Doi: 10.1186/s12894-021-00824-5.

Rahman A, Leslie SW, Desai D. Prostatic urethral lift. In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2026 Jan-. https://www.ncbi.nlm.nih.gov/books/NBK606089/. Updated August 11, 2024.

Roehrborn CG, Barkin J, Gange SN, et al. Five year results of the prospective randomized controlled prostatic urethral L.I.F.T. study. *Can J Urol*. 2017;24(3):8802-8813. https://www.canjurol.com/html/free-articles/V24I3_08_FREE_DrRoehrborn.pdf.

Rukstalis D, Grier D, Stroup SP, et al. Prostatic urethral lift (PUL) for obstructive median lobes: 12 month results of the Medlift study. *Prostate Cancer Prostatic Dis*. 2019;22(3):411-419. Doi: 10.1038/s41391-018-0118-x.

Sajan A, Mehta T, Desai P, Isaacson A, Bagla S. Minimally invasive treatments for benign prostatic hyperplasia: Systematic review and network meta-analysis. *Vasc Interv Radiol*. 2022;33(4):359-367.e8. Doi: 10.1016/j.jvir.2021.12.029.

Tanneru K, Gautam S, Norez D, et al. Meta-analysis and systematic review of intermediate-term follow-up of prostatic urethral lift for benign prostatic hyperplasia. *Int Urol Nephrol*. 2020;52(6):999-1008. Doi: 10.1007/s11255-020-02408-y.

U.S. Food and Drug Administration. De novo classification request for Neotract's UroLift System. https://www.accessdata.fda.gov/cdrh_docs/reviews/k130651.pdf. Published March 7, 2013.

U.S. Food and Drug Administration. 510(k) Premarket Notification. Implantable transprostatic tissue retractor system. https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm?ID=K173087. Published December 28, 2017.

U.S. Food and Drug Administration. 510(k) Premarket Notification database searched using product code PEW. https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm. Last updated April 14, 2025.

van Kollenburg RAA, van Riel LAMJG, de Bruin DM, de Rijke TM, Oddens JR. Novel minimally invasive treatments for lower urinary tract symptoms: a systematic review and network meta-analysis. *Int Braz J Urol*. 2023;49(4):411-427. Doi: 10.1590/S1677-5538.IBJU.2023.0016.

Xiang P, Wang M, Guan D, et al. A systematic review and meta-analysis of prostatic urethral lift for male lower urinary tract symptoms secondary to benign prostatic hyperplasia. *Eur Urol Open Sci*. 2020;19:3-15. Doi: 10.1016/j.euros.2020.05.001.

Zhang TR, Thorogood SL, Sze C. Current practice patterns in the surgical management of benign prostatic hyperplasia. *Urology*. 2023;175:157-162. Doi: 10.1016/j.urology.2023.02.025.

## Policy updates

7/2023: initial review date and clinical policy effective date: 8/2023

7/2024: Policy references updated.

8/2025: Policy references updated. Coverage modified.

8/2026: Policy references updated. Coverage modified.

## Related codes

Below are the most commonly submitted codes for the service(s)/item(s) subject to this policy CCP.1529. This is not an exhaustive list of codes. Providers are expected to consult the appropriate coding manuals and bill accordingly.

| Code | Code description |
|---|---|
| 52441 | Cystourethroscopy, with insertion of permanent adjustable transprostatic implant; single implant |
| +52442 | Cystourethroscopy, with insertion of permanent adjustable transprostatic implant; each additional permanent adjustable transprostatic implant (List separately in addition to code for primary procedure) |
| C9739 | Cystourethroscopy, with insertion of transprostatic implant; one to three implants |
| C9740 | Cystourethroscopy, with insertion of transprostatic implant; four or more implants |